

---

## EFFECT OF SHATKARMA, ASANAS AND PRANAYAMS ON HEALTH VARIABLES AMONG FEMALES

---

**Sushil**

Research Scholar,

**Dr Hari Singh**

Supervisor

Shri Krishna University, Chhatarpur (Madhya Pradesh )

---

### Abstract

The objective of the study was to assess the effect of Shatkarma, Asanas and Pranayams on selected Health Variables among females. 24 females from Spirit Yoga Centre, New Delhi randomly selected to participate in the study as subjects. The age of subjects was ranging from 25 years to 60 years. Following variables were selected for the study: Body Mass Index (BMI) and Waist-Hip ratio (WHR). The data was collected by conducting Inbody Body Composition Analyser machine and Anthropometric tape. The reliability of data was established following the instrument's reliability and tester competency. The data was collected before the commencement of training programme that was known as pre-test and thereafter, at the end of 3 months of training program known as post-test 1 and at the end of 6 months of the training programme known as post-test 2. In all, three data collection was executed from the experimental group during the 6 months specific yoga training program. A six months training program was designed to assess the effect of Shatkarma, Asanas and Pranayamas on selected variables. The training was provided to experimental group for 5 days/week for the duration of six months. Descriptive statistics and Repeated Measure MANOVA were applied using SPSS Software to analyse data. The collected data was significantly normalized before further processing. The obtained result shows the significant changes in selected variables i.e. Body Mass Index (BMI) and Waist-Hip ratio (WHR) due to specific Shatkarma, Asanas and Pranayams training.

Keywords: BMI, WHR

---

### INTRODUCTION

Regular practice of variety of Yoga techniques have been shown to lower heart rate and blood pressure in various population (Lakshmikanthan et al. 1979; Mahajan et al. 1999). Yoga exercise are scientific means for strengthen of all living or atrophying muscle fibers and tissues. This system teaches how to a make new life pulsation in active tissues. In this context it much as it is different from other system of exercise in as much as it is different from other system of exercise in as much as it teaches one how to concentrate his attention on the awakened energy which is the direct gives of power, strength and vitality to all the parts of the body. It develops the will power long with bodily strength. This aspect of yoga is technically known as a "ASANAS" which was developed by the hatha yogic into a well-organized system of physical culture. One very important part of yoga is Shatkarma

or Shatkriya as described in the yogic texts and is a very precise and systematic science. Shat means six and karma means action; shatkarma consist of six groups of purification. Main aim of shatkarmais harmony between two major pranic flows ida and pingla, it is the purification of mental , physical , balance.

With keeping the fact in mind that, the wide area of yogic exercises and its effect on various aspect of human body poorly depicted. The research scholar has decided to delimit his research work on '*Shatkarma, Asanas and Pranayams*' and its effect on selected body composition variables i.e. body weight, fat %, BMI, WHR and BMR. The following objectives were set for the present study: 1) To assess the health status of the subjects on selected variables i.e. BMI and WHR,2) To assess the effect of Shatkarma, Asanas and Pranayams on Body Mass Index (BMI) of selected female subjects and3) To assess the effect of Shatkarma, Asanas and Pranayams on Wasit-Hip Ratio (WHR) of selected female subjects.

Keeping the purpose of the study, the following hypotheses were formulated:

1. It was also hypothesized that there would be significant effect of Shatkarma, Asanas and Pranayams on Body Mass Index (BMI) of females.
2. It was also hypothesized that there would be significant effect of Shatkarma, Asanas and Pranayams on Waist-Hip Ratio (WHR) of females.

The present study was delimited as:The study was delimited to female subjects only, the study was delimited to teachers aged between 25-60 years of age and the study was further delimited to female subjects of Spirit Yoga center, New Delhi. The findings of this study had to be seen in light of some limitations as well which were as follows:The factors like diet, lifestyle, daily routine habits etc., which may have an effect on the result of the study, was considered as limitation in this study. The meteorological variations such as air temperature, atmospheric pressure, relative humidity etc. during the training period cannot be controlled and their possible influence on the study was recognized as limitation.Certain factors like past training and genetic factors that have affected the result of the study was also consider as the limitation of the study.Psychological state of the subjects may affect the result of the study. Thus, this was considered

as the limitation of study. The intent and motivation of the subjects during training and testing procedure was considered as the limitation.

Though, the present study was confined to only Shatkarma and its impact on selected Health Variables, still it may have significance in various manners. The study would profile the lifestyle and health behaviors of female subjects. The study would provide relevant information of health status of female subjects and would be eye opener for society or health experts regarding the prevailing trend of individual behaviour and its health consequences. The study would enable better policy framing health promotion measures in general. The study will serve as a motivational force to the general population to minimize the problems related to various health factors. The result of the study will be helpful for the women's which are related to the other field. The Study would provide relevant data for comparative survey of similar nature on wider population.

## **PROCEDURE AND METHODOLOGY**

### **Selection of the Subjects**

24 females from Spirit Yoga Centre, New Delhi randomly selected to participate in the study as subjects. The age of subjects was ranging from 25 years to 60 years.

### **Collection of data**

The data for the purpose of the study was collected from the female subjects of Spirit Yoga Centre, New Delhi. The data was collected before the commencement of training program that was known as pre-test and thereafter, at the end of 3 months of training program known as post-test 1 and at the end of 6 months of the training programme known as post-test 2. In all, three data collection was executed from the experimental group during the 6 months specific yoga training program.

### **Criterion Measures**

S.no	Item	Equipment/ test	Unit
1	Body Mass Index (BMI)	Body Composition Analyzer Machine	Kg/m <sup>2</sup>
2	Waist-Hip Ratio (WHR)	Anthropometric tape	cm

### **Administration of training program**

A six months training program was designed to assess the effect of Shatkarma, Asanas and Pranayamas on selected variables. The training was provided to experimental group for 5 days/week for the duration of six months. As research topic suggests, the study consists three types of exercise namely Shatkarma, Asanas and Pranayamas.

### **Statistical Techniques**

The data was collected from female subjects of Spirit Yoga Centre, New Delhi and used for the statistical treatment that specifies descriptive statistics. Various information was presented in graphical format such as Bar graph, Line graph, Pie Chart etc. when and wherever required. In order to describe the status of health condition of subjects, descriptive statistics (mean, standard deviation) was calculated. Data distribution was assessed by evaluating skewness and kurtosis along with histogram, Q-Q Plot and Box Plot diagrams. Shapiro-Wilk test was conducted for normality test and non-normality was fixed wherever required. After randomization of the data, parametric test i.e. Repeated Measure MANOVA was applied and effect of selected yogic exercises was assessed on health variables. All the statistical test was applied using SPSS (version 16) software. In all the cases of inferential statistics, 0.05 level of significance was fixed to test the hypothesis.

### **RESULT AND DISCUSSION**

The main purpose of the present study was to study the effect of selected yogic exercises i.e. Shatkarma, Asanas and Pranayams on selected health variables i.e. BMI and WHR. For the said purpose, 24 females were selected on random basis and provided Shatkarma, Asanas and Pranayams training for the period of 6 months. The minimum age of participant in the group was 26 yrs. Whereas, maximum age in the same was recorded to be 59 yrs. The mean value of age was found to be  $39.72 \pm 10.135$ . The mean value for height and weight of the same subjects were found to be  $163.42 \pm 5.02$  and  $85.596 \pm 10.68$  respectively.

**Descriptive Statistics for selected variables**

	PRE-TEST		AFTER 3 MONTHS		AFTER 6 MONTHS	
	MEAN	SD	MEAN	SD	MEAN	SD
<b>BODY MASS INDEX</b>	32.26	5.40	31.84	5.20	31.48	5.26
<b>WAIST-HIP RATIO</b>	0.99	0.06	0.96	0.05	0.94	0.05

For body mass index status of female subjects selected for the study for pre-test was recorded to be  $32.26 \pm 5.40$ . After 3 months of training, the mean value of the same was reduced to 31.84 with SD of 5.20. By the end of 6 months of specific yogic training, the mean value for the same was  $31.48 \pm 5.26$ . In case of waist hip ratio status of female subjects selected for the study for pre-test was recorded to be  $0.99 \pm 0.06$ . After 3 months of training, the mean value of the same was reduced to 0.96 with SD of 0.5. By the end of 6 months of specific yogic training, the mean value for the same was  $0.94 \pm 0.5$ .

**NORMALITY TEST**

	PRE-TEST			AFTER 3 MONTHS			AFTER 6 MONTHS		
	Statistic	df	Sig.	Statistic	df	Sig.	Statistic	df	Sig.
<b>BODY MASS INDEX</b>	.955	25	.330	.958	25	.373	.964	25	.500
<b>WAIST-HIP RATIO</b>	.964	25	.508	.947	25	.212	.956	25	.340

During the processing of data, it was found that certain variables were not normally distributed. The non-normality was assessed with the help of skewness and kurtosis data. The result was also verified with help of Shapiro-wilk test for normality along with various normality graph analysis. In case of non-normality in data distribution, it was fixed by randomization and reverified with same statistical techniques. Once the processed data fulfilled the pre-requisite assumptions, the repeated measure MANOVA technique was applied. The obtained results are shown below. Multivariate test was executed to assess the multivariate analysis of variance for all the selected variables. It is clearly shown in the table (along the "**Wilks' Lambda**" row) that  $p = .000$ . Since .000 is **less than** .05 (i.e., it satisfies  $p < .05$ ), the repeated measures MANOVA is statistically significant.

**MAUCHLEY'S TEST**

	<b>Mauchly's W</b>	<b>Approx. Chi-Square</b>	<b>df</b>	<b>Sig.</b>
<b>BODY MASS INDEX</b>	.177	38.128	2	.000
<b>WAIST-HIP RATIO</b>	.550	13.168	2	.001

Firstly, the test for sphericity was executed to assess the equality in variance in obtained data at different points of time i.e. pre-test, after 3 months of training and after 6 months of training for all the selected variables. Unfortunately, result obtained in case of all variables were found significant ( $P=0.00$ ) at 0.05 level. It validates that statistical test i.e. Repeated Measure Multivariate Analysis of Variance (MANOVA). The result found here shows that the variances of the differences between all combinations of the groups are equal. therefore, on the basis of result, we were not able to rely individually on the test above. As the sphericity of test was violated, it was corrected by making appropriate adjustment to the degree of freedom of the F-test. Thereafter, f-test and pairwise comparison were executed which has shown below.

**F-TEST**

	<b>F</b>	<b>Sig.</b>

<b>BODY MASS INDEX</b>	892.969	.000
<b>WAIST-HIP RATIO</b>	6.722E3	.000

The F-test calculation for selected variables i.e. Body Mass Index and Waist-Hip Ratio of female subjects in experimental group have shown in table above. Here, the obtained results were represented significant result for F-value as their p-value were equal to 0.00 (which is less than 0.05 level of significance) represented the significant difference in the case of all selected variables. The presented result proves the efficiency of Shatkarma, Asanas and Pranayams training in order to improve the status of body mass index and waist-hip ratio among selected subjects.

#### WITHIN SUBJECT CONTRAST TEST

	<b>Type III Sum of Squares</b>	<b>df</b>	<b>Mean Square</b>	<b>F</b>	<b>Sig.</b>
<b>BODY MASS INDEX</b>	7.434	1	7.434	8.188	.009
<b>WAIST-HIP RATIO</b>	.033	1	.033	71.319	.000

The within subject contrast test was executed to assess the trend of changes occurred in selected variables i.e. Body Mass Index and Waist-Hip Ratio due to Shatakarma, Asanas and Pranayams training program over a period of 6 months. Here, we can see within subject contrast calculations for all selected variables were found to have the linear component which was significant for main factor i.e. testing as their P-value was found less than 0.05 level of significance. On the basis of obtained result, it can be stated that 6 months of Shatakarma, Asanas and Pranayams training program might lead to linear changes in above mentioned variable i.e. Body Mass Index and Waist-Hip Ratio.

## CONCLUSIONS AND RECOMMENDATIONS

On the basis of objectives of the study and result obtained after statistical application, the following conclusions were drawn:

1. It was concluded that there is a significant effect of Shatkarma, Asanas and Pranayams on Body Mass Index (BMI) of females.
2. It was concluded that there is a significant effect of Shatkarma, Asanas and Pranayams on Waist-Hip Ratio (WHR) of females.
3. It was concluded that there is a significant linear improvement due to Shatkarma, Asanas and Pranayams on Body Mass Index (BMI) of females.
4. It was concluded that there is a significant linear improvement due to Shatkarma, Asanas and Pranayams on Waist-Hip Ratio (WHR) of females.

Above mentioned conclusions and finding has revealed many facts and filled the gap in information available regarding role of yogic exercises for betterment of health of females. Now, following recommendations are made with future research perspective:

- Similar study can be taken on male subjects as well.
- It was recommended that similar study can be carried out on profession wise as well.
- Similar study can also be conducted on a bigger population.
- A study can be conducted with including more health and body composition variables.
- Further, prediction research can be conducted to identify the health variables that should be stressed for better health condition.



- Similarly, research on factor analysis can be conducted to identify the psychological variables contributing the most to our health condition.

## References:

- Anand, B.K. (1991). Yoga and medical sciences. *Indian J PhysioPharmacol*, 35(2): 84 – 87.
- Bera, T.K., Rajapurkar, M.V., &Ganguly, S.K. (1990) Effect of yoga training on body density in school going boys. *NIS Scientific Journal*, 13,2 23-35.
- Bernardi L, et.al. (2002) Slow breathing increases arterial baroreceptor sensitivity in patients with chronic heart failure circulation, *NCBI*, 105(2), 143-5.
- Bharadwaj, I., Kulshrestha, A. &Anuja.(2013). Effect of Yogic Intervention on Blood pressure and Alpha- EEG level of working women. *Indian Journal of Traditional Knowledge*, Vol. 12 (3), pp. 542-546.
- Bhargava, R., Gogate, M.G. &Macarenhas, J.F. (1988) Autonomic responses to breath holding and its variations following pranayama, *Indian J PhysiolPharmacol*, 32(4);257–264.
- Bhutkar, P.M., et.al. (2008). Effect of Suryanamaskar Practice on Cardio-respiratory Fitness Parameters: A Pilot Study, *AJMS*, 1(2), 126-129
- Chowdhary, B. &Ghosh, B. (2015).Effect of Six Months Surya Namaskar and Selected Asanas on Body Composition Variables of Under Graduate Female Students of Jangalmahal. *Online International Interdisciplinary Research Journal*, Vol. 5(5), pp. 458-464.
- Devasena, I. &Narhare, P. (2011).Effect of yoga on heart rate and blood pressure and its clinical significance. *International Journal of Biological & Medical Research*, Vol. 2(3), pp. 750-753.
- Dhaliwal, G.S. et.al. (2014). Effects of 6-weeks Yogasana practice on physiological fitness status of university level girls. *International Journal of Physical Education, Fitness and Sports*, Vol. 3(2), pp. 43-51.
- Gadham, J., Sajja, S. &Rooha, V. (2015).Effect of yoga on obesity, hypertension and lipid profile. *International Journal of Research in Medical Sciences*, Vol. 3(5), pp. 1061-1065.
- Gore, M.M. (1984). Anatomy and physiology of yoga practices: *Ionavalakaivalayadham*, P-2.
- Karak, K., Jana, M. & Manna, A. (2015).Effect of yoga on Anthropometrical and physiological variables of college going students. *International Journal of Physical Education, Sports and Health*, Vol. 2(2), pp. 245-249.
- Khare K.C. and Kawathekar, G (2002). Lean Body Mass And Lipid Profile In Healthy Person Practicing Yoga, *Yoga Mimamsa XXXIV*. 123-128.
- Khosravi, H., Kazemzadeh, Y. &Sedaghat, S. (2015). The Effect of Yoga practice on Muscle Fitness and Body Composition in Middle age Women with Overweight. *Biological Forum – An International Journal*, Vol. 7(1), pp. 1924-1928.
- Kishore, R. & Pal, R. (2014).Effects of yogic practice in certain cardio respiratory parameters on overweight postmenopausal women. *Al Ameen Journal of Medicine Science*, Vol. 7(4), pp. 316-321.
- Lolage, R. S., &Bera, T. K. (2002).Effect of Pranayama on cardiovascularEndurance in Kho-Kho Players. *Yoga-Mimamsa*, 34(1), 13-26.
- Madhavi S. Raju, et al. (1985). Effect Of Yogic Exercises On Lean Body Mass, Skin Fold Thickness & Body Weight. *Journal of Association of Physicians of India* 33, 465-466.
- Malhotra, V. &Tandon, O. P. (2005).A study of the effect of individual Asanas on blood pressure. *Indian Journal of Traditional Knowledge*, Vol. 4(4), pp. 367-372.
- Manikandan, S. (2014). Influence of yogic practices on selected cardio respiratory system and body composition variables. *International Journal of World Research*, Vol. 1(12), pp. 14-19.
- Marger, C. F., Hicklin, L. K. & Garner, D. P. (2016). Effects of Bikram Yoga on Body Composition, Blood Pressure, and Sleep Patterns in Adult Practitioners, *Journal of Basic & Applied Sciences*, Vol. 12, pp. 75-80.
- Narayani, U. and Raj, R. L. S. P. (2010).Effect of Aerobic Training on Percentage of Body Fat, Total Cholesterol and HDL-C among Obese Women. *World Journal of Sport Sciences*, Vol. 3(1), pp. 33-36.
- Pradhan, B., andNagendra, H. (2010).Immediate effect of two yoga-basedrelaxation techniques on attention in children. *Int J Yoga*, 3(2), 67-69.

- Raja, S. C. (2014). Composition measures and high density lipoproteins among obese women. *Academic Sports Scholar*, Vol. 3(11), pp. 1-7.
- Raja, S. C. (2015). Impact of yogic practices on selected body composition measures and high density lipoproteins among obese boys. *PARIPEX-Indian Journal of Research*, Vol. 4(1), pp. 145-148.
- Rajendran, K. (2014). Effect of Yoga on Cardio Respiratory System and Body Composition of School Going Children. *International Journal of Recent Research and Applied Studies*, Vol. 1(3(20)) pp. 81-84.
- Raju P. S. et al. (1997). Influence of intensive yoga training on physiological changes in 6 adult women: a case report. *Journal of Alternative and Complementary Medicine*.3(3), 291-5.
- Shenbagavalli, A. (2005). Cardio vascular endurance and body fat percentage in relation to the practice of selected yogic exercises in students. *YogaMimamsa*, 37(1&2), 45-51.
- Udupa KN and Singh RH. (1972). The scientific basis of yoga. *J Am Med Assn*, 220(10): 1365.
- Yadav, H.K. and Singh, M.K. (2014). Effect of Surya namaskara on selected physical and physiological variables of college students. *Golden Research Thoughts*, Vol. 3(12), pp. 1-5. Yogendra, (1971). *yoga physical education*. Bombay : the yoga institution Santa, p-21.
- Zorofi, F., Hojjati, Z. & Elmiyeh, A. (2013). Effect of Yoga Exercises on the Body Composition of Fasting Females. *Journal of Fasting Health*, Vol. 1(2), pp. 70-78.